arXiv:1607.08664v2 [math.AG] 30 Nov 2016

CANONICAL SINGULARITIES OF DIMENSION THREE IN CHARACTERISTIC 2 WHICH DO NOT FOLLOW REID'S RULES

MASAYUKI HIROKADO

Abstract. We continue to study and present concrete examples in characteristic 2 of compound Du Val singularities defined over an algebraically closed field which have one dimensional singular loci but cannot be written as products (a rational double point)×(a curve) up to analytic isomorphism at any point of the loci. Unlike in other characteristics, we find a large number of such examples whose general hyperplane sections have rational double points of type D. We consider these compound Du Val singularities as a special class of canonical singularities, intend to complete classification [12, Theorem 3] in arbitrary characteristic reinforcing Miles Reid's result in characteristic zero [15, (1.14)].

.

1. Introduction

We consider varieties defined over an algebraically closed field k of characteristic $p \geq 0$. The hypersurface singularity in $\mathbf{A}^4_k$ with $p = 3$ given respectively by

$$z^2 + x^3 + y^4 + wy^3 = 0,$$
$$z^2 + x^3 + y^4 + x^2y^2 + wy^3 = 0$$

has a one dimensional singular locus C, and posesses a crepant resolution. Simple observation of blow-ups finds that these cannot be written as fiber products (a rational double point) $\times$ (a curve) up to analytic isomorphism at any $x \in C$. These are the examples of canonical singularities in positive characteristic which show that Miles Reid's result in characteristic 0 needs some modification when one generalizes it to arbitrary characteristic.

Theorem 1.1 (Reid, [15])**.** *Let X be a quasi-projective normal variety of dimension three defined over the field of complex numbers $\mathbf{C}$. If X has at most canonical singularities, then there exists some zero dimensional variety $Z \subset X$ such that for any $x \in X \setminus Z$, the complete local ring $\hat{\mathcal{O}}_{X,x}$ is either regular or isomorphic to $\mathbf{C}[[x,y,z]]/(g(x,y,z))\hat{\otimes}\mathbf{C}[[w]]$, where $\operatorname{Spec}\mathbf{C}[[x,y,z]]/(g(x,y,z))$ is a rational double point.*

One way to study such singularities is to examine versal deformations of rational double points. In the preceding work, we proved that these examples are also exhaustive in $p \geq 3$, i.e. there is no such example in $p \geq 5$ and in $p = 3$ there is no other in the following sense.

2010 *Mathematics Subject Classification.* Primary 14B05, 14G17; Secondary 14J17, 14B07.

Key words and phrases. Canonical singularities, compound Du Val singularities, rational double points, characteristic p.

Theorem 1.2 (HIS, [12]). *Let X be a three dimensional normal algebraic variety over an algebraically closed field k of characteristic $p > 0$ with at most canonical singularities. Then the following assertions hold.*

i) *X is Cohen-Macaulay in codimension two.*
ii) *Suppose $p > 2$ and that a general hyperplane section H of X has at most rational singularities. Then there exists a zero-dimensional subvariety $Z \subset X$ such that for any $x \in X \setminus Z$, the complete local ring $\hat{\mathcal{O}}_{X,x}$ is either regular or isomorphic to one of the following:*
   - *$k[[x,y,z]]/(g)\hat{\otimes}k[[w]]$ with a rational double point* $\operatorname{Spec} k[[x,y,z]]/(g)$,
   - *$k[[w,x,y,z]]/(z^2+x^3+y^4+x^2y^2+wy^3)$ with $p=3$,*
   - *$k[[w,x,y,z]]/(z^2+x^3+y^4+wy^3)$ with $p=3$.*

We call such deviations non-classical compound Du Val singularities.

TABLE 1. Non-classical compound Du Val singularities in $p = 3$.

	Type	Defining equation
E_6^1	$\mathbf{E_6^1G_2}$	$z^2+x^3+y^4+x^2y^2+wy^3=0$
E_6^0	$\mathbf{E_6^0G_2}$	$z^2+x^3+y^4+wy^3=0$

We want to complete classification. As a first step, we present in this paper concrete examples in $p = 2$.

Theorem 3.1 is our main result. See also tables 2, 3 and 4 where equations are summarized.

TABLE 2. Non-classical compound Du Val singularities in $p = 2$, II.

	Type	Defining equation
E_7^3	$\mathbf{E_7^3F_4}$	$z^2+x^3+xy^3+xyz+wy^4=0$
E_7^2	$\mathbf{E_7^2F_4}$	$z^2+x^3+xy^3+y^3z+wy^4=0$
E_7^1	$\mathbf{E_7^1F_4}$	$z^2+x^3+xy^3+wy^4=0$
E_8^3	$\mathbf{E_8^3F_4}$	$z^2+x^3+y^3z+wy^4=0$
E_8^2	$\mathbf{E_8^2F_4}$	$z^2+x^3+xy^2z+wy^4=0$
E_8^1	$\mathbf{E_8^1F_4}$	$z^2+x^3+xy^3z+wy^4=0$
E_8^0	$\mathbf{E_8^0F_4}$	$z^2+x^3+wy^4=0$

As one can see, there are quite a few compared with two equations in $p = 3$. Among them six were already known [12, Theorem 2], which are marked by boxes in the main theorem, but we believe others are new.

As for the question of rationality, it might be suggestive that if one sticks to traditional notion, these examples exhibit no peculiarity.

Corollary 4.1 *Let X be a hypersurface singularity given by one of the equations in Theorem 3.1. Then the following assertions hold.*

i) *$R^i\pi_*\mathcal{O}_{\tilde{X}} = 0$ $(i > 0)$ holds for any resolution of singularities $\pi : \tilde{X} \to X$.*
ii) *$R^i\pi_*K_{\tilde{X}} = 0$ $(i > 0)$ holds for any resolution of singularities $\pi : \tilde{X} \to X$.*

However, if one asks about F-rationality (cf. [3, §10]), these examples turn out to be highly irrational.

Proposition 4.2 *Let X be a hypersurface singularity given by one of the equations in Theorem 3.1. Then the following assertions hold.*

i) *X is F-pure if and only if the type is one of* $\mathbf{E_7^3F_4}$, $\mathbf{D_4^1B_1}$, $\mathbf{D_4^1B_2}$, $\mathbf{D_5^1C_3}$, $\mathbf{D_{2n}^{n-1}B_{n-1}}$, $\mathbf{D_{2n}^{n-1}B_n}$, $\mathbf{D_{2n}^{n-1}C_{2n-2}}$, $\mathbf{D_{2n+1}^{n-1}B_{n-1}}$, $\mathbf{D_{2n+1}^{n-1}C_{2n-1}}$ *with* $n \geq 3$.
ii) *X is not F-rational.*

2. Preliminaries

Standard notation and terminologies found for example in [11], [14] should be used without mentioning in this paper.

We distinguish rational surface singularities in the sense of Artin [1, p. 129] from those in the sense of Lipman [13, Definition (1.1)]. The former is the notion for surfaces defined over an algebraically closed field, whereas the latter is the notion for excellent surfaces.

Recent works of Cossart and Piltant [4], [5], Cutkosky [6] on resolution of singularities of three dimensional varieties in arbitrary characteristic allow us to define canonical and terminal singularities in the same way as in characteristic zero [15].

Definition 2.1 ([12])**.** Let X be a quasi-projective normal variety of dimension two or three defined over an algebraically closed field k of characteristic p. Then X is said to have only canonical (resp. terminal) singularities, if the following conditions are satisfied.

i) X is $\mathbf{Q}$-Gorenstein, i.e. there exists a positive integer m such that mK_X is a Cartier divisor.
ii) There exists a resolution of singularities $\pi : \tilde{X} \to X$, such that $mK_{\tilde{X}} \sim \pi^*(mK_X) + \sum_i a_i E_i$ with $a_i \geq 0$ (resp. $a_i > 0$), where $E := \cup_i E_i$ is the irreducible decomposition of the exceptional divisor of π.

As in characteristic zero, the definition of canonical (resp. terminal) singularities are independent of the choice of resolutions $\pi : \tilde{X} \to X$.

Proposition 2.2 (HIS, [12, Propositions 11, 12, Lemma 13])**.** *Let X be a normal variety of dimension two or three over an algebraically closed field k of characteristic $p \geq 0$, and $\pi : \tilde{X} \to X$ be a resolution, i.e. a proper birational morphism from a nonsingular $\tilde{X}$. Then for any $i > 0$, the higher direct image sheaves $R^i\pi_*\mathcal{O}_{\tilde{X}}$ and $R^i\pi_*K_{\tilde{X}}$ are independent of the choice of resolutions $\pi : \tilde{X} \to X$. In particular, if X is nonsingular, these sheaves are zero.*

3. Concrete equations

In this section we present equations of compound Du Val singularities in characteristic 2.

Theorem 3.1. *Let k be an algebraically closed field of characteristic 2, X be a hypersurface singularity defined by one of the following polynomials $f(x, y, z, w)$ as $X := \operatorname{Spec} k[x, y, z, w]/(f)$. Then the following assertions hold.*

i) *X has a one dimensional singular locus C.*
ii) *At any points $x_1, x_2 \in C$, the complete local rings $\hat{\mathcal{O}}_{X,x_1}$ and $\hat{\mathcal{O}}_{X,x_2}$ are isomorphic to each other.*
iii) *X has a crepant resolution which is obtained by a succession of blow-ups along singular loci. More detailed description is given after each equation below.*
iv) *At any point $x \in C$, the complete local ring $\hat{\mathcal{O}}_{X,x}$ cannot be expressed as $k[[x, y, z]]/(g)\hat{\otimes}k[[w]]$ with any rational double point* $\operatorname{Spec} k[[x, y, z]]/(g)$.

TABLE 3. Non-classical compound Du Val singularities in $p = 2$, I.

	Type	Defining equation
D_4^1	$\mathbf{D_4^1B_1}$	$z^2 + xyz + wx^2 + y^3 = 0$
	$\mathbf{D_4^1B_2}$	$z^2 + xy^2 + xyz + wx^2 = 0$
D_4^0	$\mathbf{D_4^0B_1}$	$z^2 + wx^2 + y^3 = 0$
	$\mathbf{D_4^0B_2}$	$z^2 + xy^2 + wx^2 = 0$
D_5^1	$\mathbf{D_5^1C_3}$	$z^2 + x^2y + y^2z + xyz + wy^3 = 0$
D_5^0	$\mathbf{D_5^0C_3}$	$z^2 + x^2y + y^2z + wy^3 = 0$
D_6^2	$\mathbf{D_6^2B_2}$	$z^2 + xy^3 + xyz + wx^2 + y^5 = 0$
	$\mathbf{D_6^2B_3}$	$z^2 + xy^3 + xyz + wx^2 = 0$
	$\mathbf{D_6^2C_4}$	$z^2 + x^2y + xy^3 + xyz + wy^4 = 0$
D_6^1	$\mathbf{D_6^1B_2}$	$z^2 + xy^3 + xy^2z + wx^2 + y^5 = 0$
	$\mathbf{D_6^1B_3}$	$z^2 + xy^3 + xy^2z + wx^2 = 0$
	$\mathbf{D_6^1C_3}$	$z^2 + x^2y + xy^3 + xy^2z + wy^3 = 0$
	$\mathbf{D_6^1C_4}$	$z^2 + x^2y + xy^3 + xy^2z + wy^4 = 0$
D_6^0	$\mathbf{D_6^0C_3}$	$z^2 + x^2y + xy^3 + wy^3 = 0$
	$\mathbf{D_6^0C_4}$	$z^2 + x^2y + xy^3 + wy^4 = 0$
D_7^2	$\mathbf{D_7^2B_2}$	$z^2 + xyz + wx^2 + y^5 = 0$
	$\mathbf{D_7^2C_5}$	$z^2 + x^2y + xyz + y^3z + wy^5 = 0$
D_7^1	$\mathbf{D_7^1C_4}$	$z^2 + x^2y + xy^2z + y^3z + wy^4 = 0$
	$\mathbf{D_7^1C_5}$	$z^2 + x^2y + xy^2z + y^3z + wy^5 = 0$
D_7^0	$\mathbf{D_7^0C_4}$	$z^2 + x^2y + y^3z + wy^4 = 0$
	$\mathbf{D_7^0C_5}$	$z^2 + x^2y + y^3z + wy^5 = 0$
D_8^3	$\mathbf{D_8^3B_3}$	$z^2 + xy^4 + xyz + wx^2 + y^7 = 0$
	$\mathbf{D_8^3B_4}$	$z^2 + xy^4 + xyz + wx^2 = 0$
	$\mathbf{D_8^3C_6}$	$z^2 + x^2y + xy^4 + xyz + wy^6 = 0$
D_8^2	$\mathbf{D_8^2B_2}$	$z^2 + xy^4 + xy^2z + wx^2 + y^5 = 0$
	$\mathbf{D_8^2B_3}$	$z^2 + xy^4 + xy^2z + wx^2 + y^7 = 0$
	$\mathbf{D_8^2B_4}$	$z^2 + xy^4 + xy^2z + wx^2 = 0$
	$\mathbf{D_8^2C_4}$	$z^2 + x^2y + xy^4 + xy^2z + wy^4 = 0$
	$\mathbf{D_8^2C_5}$	$z^2 + x^2y + xy^4 + xy^2z + wy^5 = 0$
	$\mathbf{D_8^2C_6}$	$z^2 + x^2y + xy^4 + xy^2z + wy^6 = 0$
D_8^1	$\mathbf{D_8^1B_2}$	$z^2 + xy^4 + xy^3z + wx^2 + y^5 = 0$
	$\mathbf{D_8^1B_3}$	$z^2 + xy^4 + xy^3z + wx^2 + y^7 = 0$
	$\mathbf{D_8^1B_4}$	$z^2 + xy^4 + xy^3z + wx^2 = 0$
	$\mathbf{D_8^1C_4}$	$z^2 + x^2y + xy^4 + xy^3z + wy^4 = 0$
	$\mathbf{D_8^1C_5}$	$z^2 + x^2y + xy^4 + xy^3z + wy^5 = 0$
	$\mathbf{D_8^1C_6}$	$z^2 + x^2y + xy^4 + xy^3z + wy^6 = 0$
D_8^0	$\mathbf{D_8^0B_2}$	$z^2 + xy^4 + wx^2 + y^5 = 0$
	$\mathbf{D_8^0B_3}$	$z^2 + xy^4 + wx^2 + y^7 = 0$
	$\mathbf{D_8^0B_4}$	$z^2 + xy^4 + wx^2 = 0$
	$\mathbf{D_8^0C_4}$	$z^2 + x^2y + xy^4 + wy^4 = 0$
	$\mathbf{D_8^0C_5}$	$z^2 + x^2y + xy^4 + wy^5 = 0$
	$\mathbf{D_8^0C_6}$	$z^2 + x^2y + xy^4 + wy^6 = 0$
D_9^3	$\mathbf{D_9^3B_3}$	$z^2 + xyz + wx^2 + y^7 = 0$
	$\mathbf{D_9^3C_7}$	$z^2 + x^2y + xyz + y^4z + wy^7 = 0$
D_9^2	$\mathbf{D_9^2C_5}$	$z^2 + x^2y + xy^2z + y^4z + wy^5 = 0$
	$\mathbf{D_9^2C_6}$	$z^2 + x^2y + xy^2z + y^4z + wy^6 = 0$
	$\mathbf{D_9^2C_7}$	$z^2 + x^2y + xy^2z + y^4z + wy^7 = 0$
D_9^1	$\mathbf{D_9^1C_5}$	$z^2 + x^2y + xy^3z + y^4z + wy^5 = 0$
	$\mathbf{D_9^1C_6}$	$z^2 + x^2y + xy^3z + y^4z + wy^6 = 0$
	$\mathbf{D_9^1C_7}$	$z^2 + x^2y + xy^3z + y^4z + wy^7 = 0$
D_9^0	$\mathbf{D_9^0C_5}$	$z^2 + x^2y + y^4z + wy^5 = 0$
	$\mathbf{D_9^0C_6}$	$z^2 + x^2y + y^4z + wy^6 = 0$
	$\mathbf{D_9^0C_7}$	$z^2 + x^2y + y^4z + wy^7 = 0$
$\vdots$	$\vdots$	$\vdots$

v) *A general hyperplane section $H \subset X$ has a rational double point of the type indicated on the left of each equation. For example, $\mathbf{D_4^1B_1}$ stands for that H has a rational double point of type $\mathbf{D_4^1}$ in Artin's sense, and $\mathbf{B_1}$ is the type of the rational double point in Lipman's sense which appears on H*

when one views it as a two dimensional scheme defined over the function field.

D_4^1 :
$\mathbf{D_4^1B_1}$: $z^2+xyz+wx^2+y^3=0$. *This can be resolved by a single blow-up. The exceptional divisor is irreducible and singular.*
$\mathbf{D_4^1B_2}$: $z^2+xy^2+xyz+wx^2=0$. *This can be resolved by two blow-ups. A trivial product of a rational double point of type A_1 with a curve appears after the first blow-up.*

D_4^0 :
$\boxed{\mathbf{D_4^0B_1} : z^2+wx^2+y^3=0.}$ *This can be resolved by a single blow-up. The exceptional divisor is irreducible and singular.*
$\boxed{\mathbf{D_4^0B_2} : z^2+xy^2+wx^2=0.}$ *This can be resolved by two blow-ups. After the first blow-up, a trivial product of a rational double point of type A_1 with a curve appears.*

D_5^1 :
$\mathbf{D_5^1C_3}$: $z^2+x^2y+y^2z+xyz+wy^3=0$. *This can be resolved by three blow-ups. After the first blow-up, two disjoint trivial products of A_1 with curves appear.*

D_5^0 :
$\boxed{\mathbf{D_5^0C_3} : z^2+x^2y+y^2z+wy^3=0.}$ *This can be resolved by three blow-ups. After the first blow-up, two disjoint trivial products of A_1 with curves appear.*

D_6^2 :
$\mathbf{D_6^2B_2}$: $z^2+xy^3+xyz+wx^2+y^5=0$. *This can be resolved by two blow-ups. After the first blow-up, the singularity $\mathbf{D_4^1B_1}$ appears.*
$\mathbf{D_6^2B_3}$: $z^2+xy^3+xyz+wx^2=0$. *This can be resolved by three blow-ups. After the first blow-up, the singularity of type $\mathbf{D_4^1B_2}$ appears.*
$\mathbf{D_6^2C_4}$: $z^2+x^2y+xy^3+xyz+wy^4=0$. *This can be resolved by four blow-ups. After the first blow-up, the singularity of type $\mathbf{D_4^1B_2}$ and the singularity isomorphic to a trivial product of A_1 with a curve appear.*

D_6^1 :
$\mathbf{D_6^1B_2}$: $z^2+xy^3+wx^2+y^5=0$. *This can be resolved by two blow-ups. After the first blow-up, the singularity of type $\mathbf{D_4^0B_1}$ appears.*
$\boxed{\mathbf{D_6^1B_3} : z^2+xy^3+wx^2=0.}$ *This can be resolved by three blow-ups. After the first blow-up, the singularity of type $\mathbf{D_4^0B_2}$ appears.*
$\mathbf{D_6^1C_3}$: $z^2+x^2y+xy^3+xy^2z+wy^3=0$. *This can be resolved by three blow-ups. After the first blow-up, two disjoint trivial products of A_1 with curves appear.*
$\mathbf{D_6^1C_4}$: $z^2+x^2y+xy^3+xy^2z+wy^4=0$. *This can be resolved by three blow-ups. After the first blow-up, a trivial product of A_1 with a curve and the singularity of type $\mathbf{D_4^0B_2}$ appear.*

D_6^0 :
$\mathbf{D_6^0C_3}$: $z^2+x^2y+xy^3+wy^3=0$. *This can be resolved by three blow-ups. After the first blow-up, two disjoint trivial products of A_1 with curves appear.*
$\mathbf{D_6^0C_4}$: $z^2+x^2y+xy^3+wy^4=0$. *This can be resolved by four blow-ups. After the first blow-up, the singularities which are isomorphic to a trivial product of A_1 with a curve and the one of type $\mathbf{D_4^0B_2}$ appear.*

D_7^2 :
$\mathbf{D_7^2B_2}$: $z^2+xyz+wx^2+y^5=0$. *This can be resolved by two blow-ups. After the first blow-up, the singularity of type $\mathbf{D_4^1B_1}$ appears.*

$\mathbf{D_7^2C_5}$: $z^2+x^2y+y^3z+xyz+wy^5=0$. *This can be resolved by five blow-ups. After the first blow-up, the singularities which are isomorphic to a trivial product of A_1 with a curve and the one of type* $\mathbf{D_5^1C_3}$ *appear.*

D_7^1 :
$\mathbf{D_7^1C_4}$: $z^2+x^2y+y^3z+xy^2z+wy^4=0$. *This can be resolved by four blow-ups. After the first blow-up, the singularities which are isomorphic to a trivial product of A_1 with a curve and the one of type* $\mathbf{D_4^0B_2}$ *appear.*
$\mathbf{D_7^1C_5}$: $z^2+x^2y+y^3z+xy^2z+wy^5=0$. *This can be resolved by five blow-ups. After the first blow-up, the singularities which are isomorphic to a trivial product of A_1 with a curve and the one of type* $\mathbf{D_5^0C_3}$ *appear.*

D_7^0 :
$\mathbf{D_7^0C_4}$: $z^2+x^2y+y^3z+wy^4=0$. *This can be resolved by four blow-ups. After the first blow-up, the singularities which are isomorphic to a trivial product of A_1 with a curve and the one of type* $\mathbf{D_4^0B_2}$ *appear.*
$\mathbf{D_7^0C_5}$: $z^2+x^2y+y^3z+wy^5=0$. *This can be resolved by five blow-ups. After the first blow-up, a trivial product of A_1 with a curve and the singularity of type* $\mathbf{D_5^0C_3}$ *appear.*

D_8^3 :
$\mathbf{D_8^3B_3}$: $z^2+xy^4+xyz+wx^2+y^7=0$. *This can be resolved by three blow-ups. After the first blow-up, the singularity of type* $\mathbf{D_6^2B_2}$ *appears.*
$\mathbf{D_8^3B_4}$: $z^2+xy^4+xyz+wx^2=0$. *This can be resolved by four blow-ups. After the first blow-up, the singularity of type* $\mathbf{D_6^2B_3}$ *appears.*
$\mathbf{D_8^3C_6}$: $z^2+x^2y+xy^4+xyz+wy^6=0$. *This can be resolved by six blow-ups. After the first blow-up, a trivial product of A_1 with a curve and the singularity of type* $\mathbf{D_6^2C_4}$ *appear.*

D_8^2 :
$\mathbf{D_8^2B_2}$: $z^2+xy^4+xy^2z+wx^2+y^5=0$. *This can be resolved by two blow-ups. After the first blow-up, the singularity of type* $\mathbf{D_4^0B_1}$ *appears.*
$\mathbf{D_8^2B_3}$: $z^2+xy^4+xy^2z+wx^2+y^7=0$. *This can be resolved by three blow-ups. After the first blow-up, the singularity of type* $\mathbf{D_6^1B_2}$ *appears.*
$\mathbf{D_8^2B_4}$: $z^2+xy^4+xy^2z+wx^2=0$. *This can be resolved by four blow-ups. After the first blow-up, the singularity of type* $\mathbf{D_6^1B_3}$ *appears.*
$\mathbf{D_8^2C_4}$: $z^2+x^2y+xy^4+xy^2z+wy^4=0$. *This can be resolved by four blow-ups. After the first blow-up, a trivial product of A_1 with a curve and the singularity of type* $\mathbf{D_4^0B_2}$ *appear.*
$\mathbf{D_8^2C_5}$: $z^2+x^2y+xy^4+xy^2z+wy^5=0$. *This can be resolved by five blow-ups. After the first blow-up, a trivial product of A_1 with a curve and the singularity of type* $\mathbf{D_6^1C_3}$ *appear.*
$\mathbf{D_8^2C_6}$: $z^2+x^2y+xy^4+xy^2z+wy^6=0$. *This can be resolved by six blow-ups. After the first blow-up, a trivial product of A_1 with a curve and the singularity of type* $\mathbf{D_6^1C_4}$ *appear.*

D_8^1 :
$\mathbf{D_8^1B_2}$: $z^2+xy^4+xy^3z+wx^2+y^5=0$. *This can be resolved by two blow-ups. After the first blow-up, the singularity of type* $\mathbf{D_4^0B_1}$ *appears.*
$\mathbf{D_8^1B_3}$: $z^2+xy^4+xy^3z+wx^2+y^7=0$. *This can be resolved by three blow-ups, After the first blow-up, the singularity of type* $\mathbf{D_6^1B_2}$ *appears.*
$\mathbf{D_8^1B_4}$: $z^2+xy^4+xy^3z+wx^2=0$. *This can be resolved by four blow-ups. After the first blow-up, the singularity of type* $\mathbf{D_6^1B_3}$ *appears.*
$\mathbf{D_8^1C_4}$: $z^2+x^2y+xy^4+xy^3z+wy^4=0$. *This can be resolved by four blow-ups. After the first blow-up, a trivial product of A_1 with a curve and the singularity of type* $\mathbf{D_4^0B_2}$ *appear.*

$\mathbf{D_8^1C_5}$: $z^2+x^2y+xy^4+xy^3z+wy^5=0$. *This can be resolved by five blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D_6^0C_3}$ *appear.*
$\mathbf{D_8^1C_6}$: $z^2+x^2y+xy^4+xy^3z+wy^6=0$. *This can be resolved by six blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D_6^0C_4}$ *appear.*

D_8^0 :
$\mathbf{D_8^0B_2}$: $z^2+xy^4+wx^2+y^5=0$. *This can be resolved by two blow-ups. After the first blow-up, the singularity of type* $\mathbf{D_4^0B_1}$ *appears.*
$\mathbf{D_8^0B_3}$: $z^2+xy^4+wx^2+y^7=0$. *This can be resolved by three blow-ups. Afer the first blow-up, the singularity of type* $\mathbf{D_6^1B_2}$ *appears.*
$\mathbf{D_8^0B_4}$: $z^2+xy^4+wx^2=0$. *This can be resolved by four blow-ups. After the first blow-up, the singularity of type* $\mathbf{D_6^1B_3}$ *appears.*
$\boxed{\mathbf{D_8^0C_4}: z^2+x^2y+xy^4+wy^4=0.}$ *This can be resolved by four blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D_4^0B_2}$ *appear.*
$\mathbf{D_8^0C_5}$: $z^2+x^2y+xy^4+wy^5=0$. *This can be resolved by two blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D_6^0C_3}$ *appear.*
$\mathbf{D_8^0C_6}$: $z^2+x^2y+xy^4+wy^6=0$. *This can be resolved by two blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D_6^0C_4}$ *appear.*

D_9^3 :
$\mathbf{D_9^3B_3}$: $z^2+xyz+wx^2+y^7=0$. *This can be resolved by three blow-ups. After the first blow-up, the singularity of type* $\mathbf{D_7^2B_2}$ *appears.*
$\mathbf{D_9^3C_7}$: $z^2+x^2y+y^4z+xyz+wy^7=0$. *This can be resolved by seven blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D_7^2C_5}$ *appear.*

D_9^2 :
$\mathbf{D_9^2C_5}$: $z^2+x^2y+y^4z+xy^2z+wy^5=0$. *This can be resolved by five blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D_6^1C_3}$ *appear.*
$\mathbf{D_9^2C_6}$: $z^2+x^2y+y^4z+xy^2z+wy^6=0$. *This can be resolved by six blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D_7^1C_4}$ *appear.*
$\mathbf{D_9^2C_7}$: $z^2+x^2y+y^4z+xy^2z+wy^7=0$. *This can be resolved by seven blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D_7^1C_5}$ *appear.*

D_9^1 :
$\mathbf{D_9^1C_5}$: $z^2+x^2y+y^4z+xy^3z+wy^5=0$. *This can be resolved by five blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D_6^0C_3}$ *appear.*
$\mathbf{D_9^1C_6}$: $z^2+x^2y+y^4z+xy^3z+wy^6=0$. *This can be resolved by six blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D_7^0C_4}$ *appear.*
$\mathbf{D_9^1C_7}$: $z^2+x^2y+y^4z+xy^3z+wy^7=0$. *This can be resolved by seven blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D_7^0C_5}$ *appear.*

D_9^0 :
$\mathbf{D_9^0C_5}$: $z^2+x^2y+y^4z+wy^5=0$. *This can be resolved by five blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D_6^0C_3}$ *appear.*
$\mathbf{D_9^0C_6}$: $z^2+x^2y+y^4z+wy^6=0$. *This can be resolved by six blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D_7^0C_4}$ *appear.*

$\mathbf{D_9^0C_7}$: $z^2+x^2y+y^4z+wy^7=0$. *This can be resolved by seven blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D_7^0C_5}$ *appear.*

$\vdots$

D_{2n}^{n-1} :

$\mathbf{D_{2n}^{n-1}B_{n-1}}$: $z^2+xy^n+xyz+wx^2+y^{2n-1}=0$. *This can be resolved by* $n-1$ *blow-ups. After the first blow-up, the singularity of type* $\mathbf{D_{2n-2}^{n-2}B_{n-2}}$ *appears.*

$\mathbf{D_{2n}^{n-1}B_n}$: $z^2+xy^n+xyz+wx^2=0$. *This can be resolved by* n *blow-ups. After the first blow-up, the singularity of type* $\mathbf{D_{2n-2}^{n-2}B_{n-1}}$ *appears.*

$\mathbf{D_{2n}^{n-1}C_{2n-2}}$: $z^2+x^2y+xy^n+xyz+wy^{2(n-1)}=0$. *This can be resolved by* $2n-2$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D_{2n-2}^{n-2}C_{2n-4}}$ *appear.*

$\vdots$

D_{2n}^{r} : $(n/2<r<n-1)$

$\mathbf{D_{2n}^{r}B_r}$: $z^2+xy^n+xy^{n-r}z+wx^2+y^{2r+1}=0$. *This can be resolved by* r *blow-ups. After the first blow-up, the singularity of type* $\mathbf{D_{2n-2}^{r-1}B_{r-1}}$ *appears.*

$\mathbf{D_{2n}^{r}B_{r+1}}$: $z^2+xy^n+xy^{n-r}z+y^{2r+3}+wx^2=0$. *This can be resolved by* $r+1$ *blow-ups. After the first blow-up, the singularity of type* $\mathbf{D_{2n-2}^{r-1}B_r}$ *appears.*

$\vdots$

$\mathbf{D_{2n}^{r}B_{n-1}}$: $z^2+xy^n+xy^{n-r}z+wx^2+y^{2n-1}=0$. *This can be resolved by* $n-1$ *blow-ups. After the first blow-up, the singularity of type* $\mathbf{D_{2n-2}^{r-1}B_{n-2}}$ *appears.*

$\mathbf{D_{2n}^{r}B_n}$: $z^2+xy^n+xy^{n-r}z+wx^2=0$. *This can be resolved by* n *blow-ups. After the first blow-up, the singularity of type* $\mathbf{D_{2n-2}^{r-1}B_{n-1}}$ *appears.*

$\mathbf{D_{2n}^{r}C_{2r}}$: $z^2+x^2y+xy^n+xy^{n-r}z+wy^{2r}=0$. *This can be resolved by* $2r$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D_{2n-2}^{r-1}C_{2r-2}}$ *appear.*

$\mathbf{D_{2n}^{r}C_{2r+1}}$: $z^2+x^2y+xy^n+xy^{n-r}z+wy^{2r+1}=0$. *This can be resolved by* $2r+1$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D_{2n-2}^{r-1}C_{2r-1}}$ *appear.*

$\vdots$

$\mathbf{D_{2n}^{r}C_{2n-3}}$: $z^2+x^2y+xy^n+xy^{n-r}z+wy^{2n-3}=0$. *This can be resolved by* $2n-3$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D_{2n-2}^{r-1}C_{2n-5}}$ *appear.*

$\mathbf{D_{2n}^{r}C_{2n-2}}$: $z^2+x^2y+xy^n+xy^{n-r}z+wy^{2(n-1)}=0$. *This can be resolved by* $2n-2$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D_{2n-2}^{r-1}C_{2n-4}}$ *appear.*

$\vdots$

D_{2n}^{r} : $(1\leq r\leq n/2)$

$\mathbf{D_{2n}^{r}B_{\lfloor\frac{n+1}{2}\rfloor}}$: $z^2+xy^n+xy^{n-r}z+wx^2+y^{2\lfloor\frac{n+1}{2}\rfloor+1}=0$. *This can be resolved by* $\lfloor\frac{n+1}{2}\rfloor$ *blow-ups. After the first blow-up, the singularity of type* $\mathbf{D_{2n-2}^{r-1}B_{\lfloor\frac{n-1}{2}\rfloor}}$ (*resp. of type* $\mathbf{D_{2n-4}^{\max\{0,r-2\}}B_{\lfloor\frac{n-1}{2}\rfloor}}$) *appears, if* n *is odd* (*resp.* n *is even*). *Here the symbol* $\lfloor\ \ \rfloor$ *stands for rounding down.*

$\mathbf{D_{2n}^{r}B_{\lfloor\frac{n+1}{2}\rfloor+1}}$: $z^2+xy^n+xy^{n-r}z+wx^2+y^{2\lfloor\frac{n+1}{2}\rfloor+3}=0$. *This can be resolved by* $\lfloor\frac{n+1}{2}\rfloor+1$ *blow-ups. After the first blow-up, the singularity of type* $\mathbf{D_{2n-2}^{r-1}B_{\lfloor\frac{n+1}{2}\rfloor}}$ *appears.*

$\vdots$

$\mathbf{D^r_{2n}B_{n-1}}$: $z^2+xy^n+xy^{n-r}z+wx^2+y^{2n-1}=0$. *This can be resolved by* $n-1$ *blow-ups. After the first blow-up, the singularity of type* $\mathbf{D^{r-1}_{2n-2}B_{n-2}}$ *appears.*
$\mathbf{D^r_{2n}B_n}$: $z^2+xy^n+xy^{n-r}z+wx^2=0$. *This can be resolved by* n *blow-ups. After the first blow-up, the singularity of type* $\mathbf{D^{r-1}_{2n-2}B_{n-1}}$ *appears.*
$\mathbf{D^r_{2n}C_n}$: $z^2+x^2y+xy^n+xy^{n-r}z+wy^n=0$. *This can be resolved by* n *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D^{\max\{0,\,r-2\}}_{2n-4}C_{n-2}}$ *appear.*
$\mathbf{D^r_{2n}C_{n+1}}$: $z^2+x^2y+xy^n+xy^{n-r}z+wy^{n+1}=0$. *This can be resolved by* $n+1$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D^{r-1}_{2n-2}C_{n-1}}$ *appear.*

$\vdots$

$\mathbf{D^r_{2n}C_{2n-3}}$: $z^2+x^2y+xy^n+xy^{n-r}z+wy^{2n-3}=0$. *This can be resolved by* $2n-3$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D^{r-1}_{2n-2}C_{2n-5}}$ *appear.*
$\mathbf{D^r_{2n}C_{2n-2}}$: $z^2+x^2y+xy^n+xy^{n-r}z+wy^{2n-2}=0$. *This can be resolved by* $2n-2$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D^{r-1}_{2n-2}C_{2n-4}}$ *appear.*

$\vdots$

D^0_{2n} :
$\mathbf{D^0_{2n}B_{n/2}}$: $z^2+xy^n+wx^2+y^{n+1}=0$ *with* n *even. This can be resolved by* $n/2$ *blow-ups. After the first blow-up, the singularity of type* $\mathbf{D^0_{2n-4}B_{(n-2)/2}}$ *appears.*
$\mathbf{D^0_{2n}B_{(n+2)/2}}$: $z^2+xy^n+wx^2+y^{n+3}=0$ *with* n *even. This can be resolved by* $(n+2)/2$ *blow-ups. After the first blow-up, the singularity of type* $\mathbf{D^1_{2n-2}B_{n/2}}$ *appears.*

$\vdots$

$\mathbf{D^0_{2n}B_{n-1}}$: $z^2+xy^n+wx^2+y^{2n-1}=0$ *with* n *even. This can be resolved by* $n-1$ *blow-ups. After the first blow-up, the singularity of type* $\mathbf{D^1_{2n-2}B_{n-2}}$ *appears.*
$\mathbf{D^0_{2n}B_n}$: $z^2+xy^n+wx^2=0$ *with* n *even. This can be resolved by* n *blow-ups. After the first blow-up, the singularity of type* $\mathbf{D^1_{2n-2}B_{n-1}}$ *appears.*
$\mathbf{D^0_{2n}C_n}$: $z^2+x^2y+wy^n=0$. *This can be resolved by* n *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D^0_{2n-4}C_{n-2}}$ *appear.*
$\mathbf{D^0_{2n}C_{n+1}}$: $z^2+x^2y+xy^n+wy^{n+1}=0$. *This can be resolved by* $n+1$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D^0_{2n-2}C_{n-1}}$ *appear.*

$\vdots$

$\mathbf{D^0_{2n}C_{2n-3}}$: $z^2+x^2y+xy^n+wy^{2n-3}=0$. *This can be resolved by* $2n-3$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D^0_{2n-2}C_{2n-5}}$ *appear.*
$\mathbf{D^0_{2n}C_{2n-2}}$: $z^2+x^2y+xy^n+wy^{2(n-1)}=0$. *This can be resolved by* $2n-2$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D^0_{2n-2}C_{2n-4}}$ *appear.*

D^{n-1}_{2n+1} :
$\mathbf{D^{n-1}_{2n+1}B_{n-1}}$: $z^2+xyz+wx^2+y^{2n-1}=0$. *This can be resolved by* $n-1$ *blow-ups. After the first blow-up, the singularity of type* $\mathbf{D^{n-2}_{2n-1}B_{n-2}}$ *appears.*
$\mathbf{D^{n-1}_{2n+1}C_{2n-1}}$: $z^2+x^2y+y^nz+xyz+wy^{2n-1}=0$. *This can be resolved by* $2n-1$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D^{n-2}_{2n-1}C_{2n-3}}$ *appear.*

D^r_{2n+1} : $(n/2 < r < n-1)$.

$\mathbf{D^r_{2n+1}C_{2r+1}}$: $z^2 + x^2y + y^nz + xy^{n-r}z + wy^{2r+1} = 0$. *This can be resolved by* $2r+1$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D^{r-1}_{2n-1}C_{2r-1}}$ *appear.*

$\vdots$

$\mathbf{D^r_{2n+1}C_{2n-2}}$: $z^2 + x^2y + y^nz + xy^{n-r}z + wy^{2n-2} = 0$. *This can be resolved by* $2n-2$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D^{r-1}_{2n-1}C_{2n-4}}$ *appear.*

$\mathbf{D^r_{2n+1}C_{2n-1}}$: $z^2 + x^2y + y^nz + xy^{n-r}z + wy^{2n-1} = 0$. *This can be resolved by* $2n-1$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D^{r-1}_{2n-1}C_{2n-3}}$ *appear.*

D^r_{2n+1} : $(1 \le r \le n/2)$.

$\mathbf{D^r_{2n+1}C_{n+1}}$: $z^2 + x^2y + y^nz + xy^{n-r}z + wy^{n+1} = 0$. *This can be resolved by* $n+1$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D^{r-1}_{2n-2}C_{n-1}}$ *appear.*

$\mathbf{D^r_{2n+1}C_{n+2}}$: $z^2 + x^2y + y^nz + xy^{n-r}z + wy^{n+2} = 0$. *This can be resolved by* $n+2$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D^{r-1}_{2n-1}C_n}$ *appear.*

$\vdots$

$\mathbf{D^r_{2n+1}C_{2n-2}}$: $z^2 + x^2y + y^nz + xy^{n-r}z + wy^{2n-2} = 0$. *This can be resolved by* $2n-2$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D^{r-1}_{2n-1}C_{2n-4}}$ *appear.*

$\mathbf{D^r_{2n+1}C_{2n-1}}$: $z^2 + x^2y + y^nz + xy^{n-r}z + wy^{2n-1} = 0$. *This can be resolved by* $2n-1$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D^{r-1}_{2n-1}C_{2n-3}}$ *appear.*

D^0_{2n+1} :

$\mathbf{D^0_{2n+1}C_{n+1}}$: $z^2 + x^2y + y^nz + wy^{n+1} = 0$. *This can be resolved by* $n+1$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D^0_{2n-2}C_{n-1}}$ *appear.*

$\mathbf{D^0_{2n+1}C_{n+2}}$: $z^2 + x^2y + y^nz + wy^{n+2} = 0$. *This can be resolved by* $n+2$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D^0_{2n-1}C_n}$ *appear.*

$\vdots$

$\mathbf{D^0_{2n+1}C_{2n-2}}$: $z^2 + x^2y + y^nz + wy^{2n-2} = 0$. *This can be resolved by* $2n-2$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D^0_{2n-1}C_{2n-4}}$ *appear.*

$\mathbf{D^0_{2n+1}C_{2n-1}}$: $z^2 + x^2y + y^nz + wy^{2n-1} = 0$. *This can be resolved by* $2n-1$ *blow-ups. After the first blow-up, a trivial product of* A_1 *with a curve and the singularity of type* $\mathbf{D^0_{2n-1}C_{2n-3}}$ *appear.*

E^3_7 :

$\mathbf{E^3_7F_4}$: $z^2 + x^3 + xy^3 + xyz + wy^4 = 0$. *This can be resolved by four blow-ups. After the first blow-up, the singularity of type* $\mathbf{D^2_6B_3}$ *appears.*

E^2_7 :

$\mathbf{E^2_7F_4}$: $z^2 + x^3 + xy^3 + y^3z + wy^4 = 0$. *This can be resolved by three blow-ups. After the first blow-up, the singularity of type* $\mathbf{D^1_6B_3}$ *appears.*

E^1_7 :

$\mathbf{E^1_7F_4}$: $z^2 + x^3 + xy^3 + wy^4 = 0$. *This can be resolved by four blow-ups. After the first blow-up, the singularity of type* $\mathbf{D^1_6B_3}$ *appears.*

E_8^3 :
$\mathbf{E_8^3F_4}$: $z^2+x^3+y^3z+wy^4=0$. *This can be resolved by four blow-ups. After the first blow-up, the singularity of type* $\mathbf{D_6^1B_3}$ *appears.*
E_8^2 :
$\mathbf{E_8^2F_4}$: $z^2+x^3+xy^2z+wy^4=0$. *This can be resolved by three blow-ups. After the first blow-up, the singularity of type* $\mathbf{D_6^1B_3}$ *appears.*

E_8^1 :
$\mathbf{E_8^1F_4}$: $z^2+x^3+xy^3z+wy^4=0$. *This can be resolved by four blow-ups. After the first blow-up, the singularity of type* $\mathbf{D_6^1B_3}$ *appears.*
E_8^0 :
$\boxed{\mathbf{E_8^0F_4} : z^2+x^3+wy^4=0.}$ *This can be resolved by four blow-ups. After the first blow-up, the singularity of type* $\mathbf{D_6^1B_3}$ *appears.*

Proof. We divide the set of equations in question into three subsets, i.e. the subset consisting of equations labeled as type D_4^1 through D_7^0 (cf. Table 3), the subset consisting of equations of type D_{2n}^{n-1} through D_{2n+1}^0 with $n \geq 4$ (cf. Table 4) and thirdly the subset consisting of equations labeled as type E_7^3 through E_8^0 (cf. Table 2). We prove the assertion for type D_N by induction on $N \geq 4$, then one by one for type E_7, E_8.

As the proof of i), ii), iii) and iv) consists of repetition of elementary procedures, i.e. blowing up, applying Jacobian criterion for regularity, exchanging coordinates, etc., we present here only the essential step of induction for assertion iii) and iv), i.e. type D_{2n}^{n-1} through D_{2n+1}^0 with $n \geq 4$.

By Jacobian criterion for regularity over an algebraically closed field k, we find that the line defined by $x=y=z=0$ is the singular locus of each hypersurface. So we blow up this line and examine the resulting singularities first in the chart $x'=x/y$, $y'=y$, $z'=z/y$, (w remains unchanged). The equations are obtained easily (cf. the tables below which should be combined with Table 4), so we identify the equations using the induction hypothesis. Verification goes straightforward, however some coordinate changes are required in the following cases.

For the singularity of type $\mathbf{D_{2n}^r B_{\lfloor (n+1)/2 \rfloor}}$ with $1 \leq r \leq n/2$ and $4 \leq n$, we have the equation after a blow-up $(z')^2+x'(y')^{n-1}+x'(y')^{n-r}z'+w(x')^2+(y')^{2\lfloor \frac{n+1}{2} \rfloor -1}=0$. First we consider the case n is even. Introduce a new coordinate $y'':=y'+x'$, then the equation is $(z')^2+x'(x'+y'')^{n-1}+x'(x'+y'')^{n-r}z'+w(x')^2+(x'+y'')^{n-1}=0$. Then factor out $(x')^2$ and obtain $(z')^2+x'(y'')^{n-r}z'+x'(y'')^{n-2}+u_1(y'')^{n-1}+w'(x')^2=0$ with u_1 a unit in $k[[x',y'',z',w]]$ and an appropriate coordinate w'. If $r>2$, we multiply x',y'',z',w' by some units and may replace u_1 by 1. The resulting singularity is of type $\mathbf{D_{2(n-2)}^{r-2}B_{\lfloor \frac{(n-2)+1}{2} \rfloor}}$. If $r \leq 2$, we put $u_2 := 1+(y'')^{2-r}z'$ and have the equation $(z')^2+u_2x'(y'')^{n-2}+u_1(y'')^{n-1}+w'(x')^2=0$ with u_1, u_2 units. We may replace u_1, u_2 by 1 in a similar way as above. This is the singularity of type $\mathbf{D_{2(n-2)}^0 B_{(n-2)/2}}$. Secondly we consider the case n is odd. But a straightforward checking finds the singularity is of type $\mathbf{D_{2(n-1)}^{r-1}B_{\lfloor \frac{(n-1)+1}{2} \rfloor}}$.

For the singularity of type $\mathbf{D_{2n}^r C_n}$ with $1 \leq r \leq n/2$ and $4 \leq n$, after a blow-up, we have the hypersurface $(z')^2+(x')^2y'+x'(y')^{n-1}+x'(y')^{n-r}z'+w(y')^{n-2}=0$. If $r>2$, this is the singularity of type $\mathbf{D_{2(n-2)}^{r-2}C_{n-2}}$. If $r \leq 2$, we introduce a new coordinate $w' := w+x'(y')^{2-r}z'$ and have $(z')^2+(x')^2y'+x'(y')^{n-1}+w'(y')^{n-2}=0$. This is the singularity of type $\mathbf{D_{2(n-2)}^0 C_{n-2}}$ if $n \geq 5$, and of type $\mathbf{D_4^0 B_2}$ if $n=4$.

For $\mathbf{D^0_{2n}B_{n/2}}$ with $4 \leq n$ even, the hypersurface after a blow-up is $(z')^2 + x'(y')^{n-1} + y'^{n-1} + w(x')^2 = 0$. Another new coordinate $y'' := y' + x'$ gives $(z')^2 + x'(y'' + x')^{n-1} + (y'' + x')^{n-1} + w(x')^2 = 0$. Then factoring out $(x')^2$, we have $(z')^2 + u_1 x'(y'')^{n-2} + (y'')^{n-1} + w'(x')^2 = 0$ with an appropriate coordinate w' and a unit u_1. This u_1 may be replaced by 1 and the singularity is of type $\mathbf{D^0_{2(n-2)}B_{(n-2)/2}}$.

	Original	Equation after a blow-up $x' = x/y$, $y' = y$, $z' = z/y$.	Resulting
D^r_{2n}	$\mathbf{D^r_{2n}B_r}$	$y'^2[z'^2 + x'y'^{n-1} + x'y'^{n-r}z' + wx'^2 + y'^{2r-1}] = 0$	$\mathbf{D^{r-1}_{2n-2}B_{r-1}}$
	$\mathbf{D^r_{2n}B_{r+1}}$	$y'^2[z'^2 + x'y'^{n-1} + x'y'^{n-r}z' + wx'^2 + y'^{2r+1}] = 0$	$\mathbf{D^{r-1}_{2n-2}B_r}$
	⋮	⋮	⋮
	$\mathbf{D^r_{2n}B_{n-1}}$	$y'^2[z'^2 + x'y'^{n-1} + x'y'^{n-r}z' + wx'^2 + y'^{2n-3}] = 0$	$\mathbf{D^{r-1}_{2n-2}B_{n-2}}$
	$\mathbf{D^r_{2n}B_n}$	$y'^2[z'^2 + x'y'^{n-1} + x'y'^{n-r}z' + wx'^2] = 0$	$\mathbf{D^{r-1}_{2n-2}B_{n-1}}$
	$\mathbf{D^r_{2n}C_{2r}}$	$y'^2[z'^2 + x'^2y' + x'y'^{n-1} + x'y'^{n-r}z' + wy'^{2r-2}] = 0$	$\mathbf{D^{r-1}_{2n-2}C_{2r-2}}$
	$\mathbf{D^r_{2n}C_{2r+1}}$	$y'^2[z'^2 + x'^2y' + x'y'^{n-1} + x'y'^{n-r}z' + wy'^{2r-1}] = 0$	$\mathbf{D^{r-1}_{2n-2}C_{2r-1}}$
	⋮	⋮	⋮
	$\mathbf{D^r_{2n}C_{2n-2}}$	$y'^2[z'^2 + x'^2y' + x'y'^{n-1} + x'y'^{n-r}z' + wy'^{2n-4}] = 0$	$\mathbf{D^{r-1}_{2n-2}C_{2n-4}}$
⋮	⋮	⋮	
D^r_{2n}	$\mathbf{D^r_{2n}B_{\lfloor (n+1)/2 \rfloor}}$	$y'^2[z'^2 + x'y'^{n-1} + x'y'^{n-r}z' + wx'^2 + y'^{2\lfloor (n+1)/2 \rfloor - 1}] = 0$	(*)
	$\mathbf{D^r_{2n}B_{\lfloor (n+1)/2 \rfloor + 1}}$	$y'^2[z'^2 + x'y'^{n-1} + x'y'^{n-r}z' + wx'^2 + y'^{2\lfloor (n+1)/2 \rfloor + 1}] = 0$	(**)
	⋮	⋮	⋮
	$\mathbf{D^r_{2n}B_{n-1}}$	$y'^2[z'^2 + x'y'^{n-1} + x'y'^{n-r}z' + wx'^2 + y'^{2n-3}] = 0$	$\mathbf{D^{r-1}_{2n-2}B_{n-2}}$
	$\mathbf{D^r_{2n}B_n}$	$y'^2[z'^2 + x'y'^{n-1} + x'y'^{n-r}z' + wx'^2] = 0$	$\mathbf{D^{r-1}_{2n-2}B_{n-1}}$
	$\mathbf{D^r_{2n}C_n}$	$y'^2[z'^2 + x'^2y' + x'y'^{n-1} + x'y'^{n-r}z' + wy'^{n-2}] = 0$	(***)
	$\mathbf{D^r_{2n}C_{n+1}}$	$y'^2[z'^2 + x'^2y' + x'y'^{n-1} + x'y'^{n-r}z' + wy'^{n-1}] = 0$	$\mathbf{D^{r-1}_{2(n-1)}C_{n-1}}$
	⋮	⋮	⋮
	$\mathbf{D^r_{2n}C_{2n-3}}$	$y'^2[z'^2 + x'^2y' + x'y'^{n-1} + x'y'^{n-r}z' + wy'^{2(n-1)-3}] = 0$	$\mathbf{D^{r-1}_{2(n-1)}C_{2n-5}}$
	$\mathbf{D^r_{2n}C_{2n-2}}$	$y'^2[z'^2 + x'^2y' + x'y'^{n-1} + x'y'^{n-r}z' + wy'^{2(n-1)-2}] = 0$	$\mathbf{D^{r-1}_{2(n-1)}C_{2n-4}}$
⋮	⋮	⋮	
D^0_{2n}	$\mathbf{D^0_{2n}B_{n/2}}$	$y'^2[z'^2 + x'y'^{n-1} + wx'^2 + y'^{n-1}] = 0$	$\mathbf{D^0_{2n-4}B_{(n-2)/2}}$
	$\mathbf{D^0_{2n}B_{(n+2)/2}}$	$y'^2[z'^2 + x'y'^{n-1} + wx'^2 + y'^{n+1}] = 0$	$\mathbf{D^1_{2n-2}B_{n/2}}$
	⋮	⋮	⋮
	$\mathbf{D^0_{2n}B_{n-1}}$	$y'^2[z'^2 + x'y'^{n-1} + wx'^2 + y'^{2n-3}] = 0$	$\mathbf{D^1_{2n-2}B_{n-2}}$
	$\mathbf{D^0_{2n}B_n}$	$y'^2[z'^2 + x'y'^{n-1} + wx'^2] = 0$	$\mathbf{D^1_{2n-2}B_{n-1}}$
	$\mathbf{D^0_{2n}C_n}$	$y'^2[z'^2 + x'^2y' + x'y'^{n-1} + wy'^{n-2}] = 0$	$\mathbf{D^0_{2n-4}C_{n-2}}$
	$\mathbf{D^0_{2n}C_{n+1}}$	$y'^2[z'^2 + x'^2y' + x'y'^{n-1} + wy'^{n-1}] = 0$	$\mathbf{D^0_{2n-2}C_{n-1}}$
	⋮	⋮	⋮
	$\mathbf{D^0_{2n}C_{2n-3}}$	$y'^2[z'^2 + x'^2y' + x'y'^{n-1} + wy'^{2n-5}] = 0$	$\mathbf{D^0_{2n-2}C_{2n-5}}$
	$\mathbf{D^0_{2n}C_{2n-2}}$	$y'^2[z'^2 + x'^2y' + x'y'^{n-1} + wy'^{2n-4}] = 0$	$\mathbf{D^0_{2n-2}C_{2n-4}}$

(*) $\mathbf{D^{\max\{0,r-2\}}_{2(n-2)}B_{\lfloor \frac{n-1}{2} \rfloor}}$ if n is even, $\mathbf{D^{r-1}_{2(n-1)}B_{\lfloor \frac{n-1}{2} \rfloor}}$ if n is odd.

(**) $\mathbf{D^1_{2(n-1)}B_{n-1}}$ if n is even and $r = 1$, $\mathbf{D^{r-1}_{2(n-1)}B_{\lfloor \frac{n+1}{2} \rfloor}}$ otherwise.

(***) $\mathbf{D^{\max\{0,r-2\}}_{2(n-2)}C_{n-2}}$ if $n \geq 5$, $\mathbf{D^0_4B_2}$ if $n = 4$.

For $\mathbf{D^0_{2n}B_{(n+2)/2}}$ with $4 \leq n$ even, we have $(z')^2 + x'(y')^{n-1} + (y')^{n+1} + w(x')^2 = 0$ after a blow-up. Then introduce a new coordinate $y'' := y' + z'$, we have $(z')^2 + x'(y'' + z')^{n-1} + (y'' + z')^{n+1} + w(x')^2 = 0$. Then factor out $(z')^2$ and we have $u_1(z')^2 + x'(y'')^{n-1} + x'(y'')^{n-2}z' + (y'')^{n+1} + (y'')^n z' + w(x')^2 = 0$ with a unit u_1 in $k[[x', y'', z', w]]$. Another new coordinate $x'' := x' + y''z'$ makes the equation $u_2(z')^2 + x''(y'')^{n-1} + x''(y'')^{n-2}z' + (y'')^{n+1} + w(x'')^2 = 0$ with a unit u_2. This unit

u_2 may be replaced by 1 by multiplying x'', y'', z', w by some units and we have the singularity of type $\mathbf{D^1_{2(n-1)}B_{n/2}}$.

$\vdots$

For $\mathbf{D^0_{2n}B_{n-1}}$ with $4 \leq n$ even, we have $(z')^2 + x'(y')^{n-1} + (y')^{2n-3} + w(x')^2 = 0$ after a blow-up. Then introduce new coordinates as in $\mathbf{D^0_{2n}B_{(n+2)/2}}$ above, we have $u_2(z')^2 + x''(y'')^{n-1} + x''(y'')^{n-2}z' + (y'')^{2n-3} + w(x'')^2 = 0$ with a unit u_2. Multiplying x'', y'', z', w by some units, we replace u_2 by 1 and have the singularity of type $\mathbf{D^1_{2(n-1)}B_{(n-1)-1}}$.

For $\mathbf{D^0_{2n}B_n}$ with $4 \leq n$ even, we have $(z')^2 + x'(y')^{n-1} + w(x')^2 = 0$ after a blow-up. Then introduce a new coordinate $y'' := y' + z'$, we have $z'^2 + x'(y'' + z')^{n-1} + w(x')^2 = 0$. Then factoring out $(z')^2$, we have $u_1(z')^2 + x'(y'')^{n-1} + x'(y'')^{n-2}z' + w(x')^2 = 0$ with a unit u_1. Then multiplying x', y'', z', w by some units again, we may replace u_1 by 1 and have the singularity of type $\mathbf{D^1_{2(n-1)}B_{n-1}}$.

For $\mathbf{D^0_{2n}C_n}$ with $4 \leq n$, we have $(z')^2 + (x')^2y' + x'(y')^{n-1} + w(y')^{n-2} = 0$ after a blow-up. A new coordinate $w' := w + x'y' + x'$ makes the equation $(z')^2 + (x')^2y' + x'(y')^{n-2} + w'(y')^{n-2} = 0$. This is the singularity of type $\mathbf{D^0_{2(n-2)}C_{n-2}}$.

	Original	Equation after a blow-up $x' = x/y$, $y' = y$, $z' = z/y$.	Resulting
D^{n-1}_{2n+1}	$\mathbf{D^{n-1}_{2n+1}B_{n-1}}$	$y'^2[z'^2 + x'y'z' + wx'^2 + y'^{2n-3}] = 0$	$\mathbf{D^{n-2}_{2n-1}B_{n-2}}$
	$\mathbf{D^{n-1}_{2n+1}C_{2n-1}}$	$y'^2[z'^2 + x'^2y' + y'^{n-1}z' + x'y'z' + wy'^{2n-3}] = 0$	$\mathbf{D^{n-2}_{2n-1}C_{2n-3}}$
$\vdots$	$\vdots$	$\vdots$	
D^r_{2n+1}	$\mathbf{D^r_{2n+1}C_{2r+1}}$	$y'^2[z'^2 + x'^2y' + x'y'^{n-r}z' + y'^{n-1}z' + wy'^{2r-1}] = 0$	$\mathbf{D^{r-1}_{2n-1}C_{2r-1}}$
	$\mathbf{D^r_{2n+1}C_{2r+2}}$	$y'^2[z'^2 + x'^2y' + x'y'^{n-r}z' + y'^{n-1}z' + wy'^{2r}] = 0$	$\mathbf{D^{r-1}_{2n-1}C_{2r}}$
	$\vdots$	$\vdots$	$\vdots$
	$\mathbf{D^r_{2n+1}C_{2n-2}}$	$y'^2[z'^2 + x'^2y' + x'y'^{n-r}z' + y'^{n-1}z' + wy'^{2n-4}] = 0$	$\mathbf{D^{r-1}_{2n-1}C_{2n-4}}$
	$\mathbf{D^r_{2n+1}C_{2n-1}}$	$y'^2[z'^2 + x'^2y' + x'y'^{n-r}z' + y'^{n-1}z' + wy'^{2n-3}] = 0$	$\mathbf{D^{r-1}_{2n-1}C_{2n-3}}$
$\vdots$	$\vdots$	$\vdots$	
D^r_{2n+1}	$\mathbf{D^r_{2n+1}C_{n+1}}$	$y'^2[z'^2 + x'^2y' + x'y'^{n-r}z' + y'^{n-1}z' + wy'^{n-1}] = 0$	$\mathbf{D^{r-1}_{2n-2}C_{n-1}}$
	$\mathbf{D^r_{2n+1}C_{n+2}}$	$y'^2[z'^2 + x'^2y' + x'y'^{n-r}z' + y'^{n-1}z' + wy'^{n}] = 0$	$\mathbf{D^{r-1}_{2n-1}C_{n}}$
	$\vdots$	$\vdots$	$\vdots$
	$\mathbf{D^r_{2n+1}C_{2n-2}}$	$y'^2[z'^2 + x'^2y' + x'y'^{n-r}z' + y'^{n-1}z' + wy'^{2n-4}] = 0$	$\mathbf{D^{r-1}_{2n-1}C_{2n-4}}$
	$\mathbf{D^r_{2n+1}C_{2n-1}}$	$y'^2[z'^2 + x'^2y' + x'y'^{n-r}z' + y'^{n-1}z' + wy'^{2n-3}] = 0$	$\mathbf{D^{r-1}_{2n-1}C_{2n-3}}$
$\vdots$	$\vdots$	$\vdots$	
D^0_{2n+1}	$\mathbf{D^0_{2n+1}C_{n+1}}$	$y'^2[z'^2 + x'^2y' + y'^{n-1}z' + wy'^{n-1}] = 0$	$\mathbf{D^0_{2n-2}C_{n-1}}$
	$\mathbf{D^0_{2n+1}C_{n+2}}$	$y'^2[z'^2 + x'^2y' + y'^{n-1}z' + wy'^{n}] = 0$	$\mathbf{D^0_{2n-1}C_{n}}$
	$\vdots$	$\vdots$	$\vdots$
	$\mathbf{D^0_{2n+1}C_{2n-2}}$	$y'^2[z'^2 + x'^2y' + y'^{n-1}z' + wy'^{2n-4}] = 0$	$\mathbf{D^0_{2n-1}C_{2n-4}}$
	$\mathbf{D^0_{2n+1}C_{2n-1}}$	$y'^2[z'^2 + x'^2y' + y'^{n-1}z' + wy'^{2n-3}] = 0$	$\mathbf{D^0_{2n-1}C_{2n-3}}$

For $\mathbf{D^r_{2n+1}C_{n+1}}$ with $1 \leq r \leq n/2$ and $4 \leq n$. We have $(z')^2 + (x')^2y' + x'(y')^{n-r}z' + (y')^{n-1}z + w(y')^{n-1} = 0$. A new coordinate $w' := w + z'$ makes the equation $(z')^2 + (x')^2y' + x'(y')^{n-r}z' + w(y')^{n-1} = 0$. This is the singularity of type $\mathbf{D^{r-1}_{2(n-1)}C_{n-1}}$.

For the singularity of type $\mathbf{D^0_{2n+1}C_{n+1}}$ with $1 \leq r \leq n/2$ and $4 \leq n$. We have $(z')^2 + (x')^2y' + (y')^{n-1}z' + w(y')^{n-1} = 0$. A new coordinate $w' := w + z'$ makes the equation $(z')^2 + (x')^2y' + w'(y')^{n-1} = 0$. This is the singularity of type $\mathbf{D^0_{2(n-1)}C_{n-1}}$.

In the other chart $x'' = x$, $y'' = y/x$, $z'' = z/x$, (w remains unchanged), for the equations labeled with $\mathbf{C_n}$ we find a trivial product of the rational double point of type A_1 and a nonsingular curve, but for other equations the resulting surfaces are nonsingular. This makes the induction work, and we obtain assertions iii) and iv).

For the assertion v), we consider a generic hyperplane section and substitute w by $\xi x + \eta y + \theta z + \iota$ with variables ξ, η, θ, ι. Each polynomial listed in Tables 2, 3, 4 is viewed as an element of $k(\xi, \eta, \theta, \iota)[x, y, z]$. Then we apply the classification algorithm of Lipman [13, §24] first over the function field $k(\xi, \eta, \theta, \iota)$, then we examine how the field extension of $k(\xi, \eta, \theta, \iota)$ affects the singularities. If the invariant τ of the quadratic form $Q(X, Y, Z)$ (loc. cit.) satisfies $\tau \geq 2$, we have a generic hyperplane section with the singularity of type B_n. If $\tau = 1$, then we consider the cubic form $\overline{\mathrm{G}}(\mathrm{U}, \mathrm{V}) \in k(\xi, \eta, \theta, \iota)[U, V]$ (loc. cit.). If $\overline{\mathrm{G}}(\mathrm{U}, \mathrm{V})$ is the product of a linear and an irreducible quadratic factor over $k(\xi, \eta, \theta, \iota)$, we have the singularity of type C_3 . If $\overline{\mathrm{G}}(\mathrm{U}, \mathrm{V}) = \mathrm{UV}^2$, we have type C_n with $n \geq 4$, and if $\overline{\mathrm{G}}(\mathrm{U}, \mathrm{V}) = a\mathrm{V}^3$ ($a \neq 0$), we have type F_4. For singularities over an algebraically closed field $\overline{k(\xi, \eta, \theta, \iota)}$, we need examine if the quadratic form $Q(X, Y, Z)$ as well as the cubic form $\overline{\mathrm{G}}(\mathrm{U}, \mathrm{V})$ are factorized further. □

Remarks 3.2. i) Note that when we consider a three dimensional singularity, taking general hyperplane sections and blowing up singular loci are not compatible in general.

ii) We substitute w with $\xi x+\eta y+\theta z+\iota$ and consider each equation $f(x, y, z, \xi z+\eta y+\theta z+\iota) = 0$ given by an element of $k(\xi, \eta, \theta, \iota)[x, y, z]$. Then the Jacobian ideal is given by

$$(\partial F/\partial x, \partial F/\partial y, \partial F/\partial z, F) \subset k(\xi, \eta, \theta, \iota)[x, y, z]$$

with $F := f(x, y, z, \xi z + \eta y + \theta z + \iota)$ and the derivations are considered as elements of $\mathrm{Der}_{k(\xi,\eta,\theta,\iota)}(k(\xi, \eta, \theta, \iota)[x, y, z])$. We calculate its Gröbner bases. The result is shown below. (We used the lexicographical monomial order with $y > z > x$ for type $\mathbf{D^r_N B}_-$, with $x > z > y$ for type $\mathbf{D^r_N C}_-$ and with $x > y > z$ for type $\mathbf{E^r_N F_4}$. For type $\mathbf{D^r_N}$, we present here up to $\mathbf{D^0_{11} B_4}$ and $\mathbf{D^0_{11} C_9}$.)

$\mathbf{D^1_4 B_1} : y^2 + zx + \eta x^2, yz + \xi x^2, yx + \theta x^2, z^2 + \iota x^2, zx^2, x^3.$
$\mathbf{D^1_4 B_2} : y^2 + yz + \xi x^2, yx + \theta x^2, z^2 + \iota x^2, zx + \eta x^2, x^3.$
$\mathbf{D^0_4 B_1} : y^2, z^2, x^2.$
$\mathbf{D^0_4 B_2} : y^2, z^2, x^2.$
$\mathbf{D^2_6 B_2} : y^3 + yz + \xi x^2, y^2 z + zx + \eta x^2, yx + \theta x^2, z^2 + \iota x^2, zx^2, x^3.$
$\mathbf{D^2_6 B_3} : y^3 + yz + \xi x^2, yx + \theta x^2, z^2 + \iota x^2, zx + \eta x^2, x^3.$
$\mathbf{D^1_6 B_2} : y^3 + y^2 z, y^2 x, z^2, x^2.$
$\mathbf{D^1_6 B_3} : y^3 + y^2 z, y^2 x, z^2, x^2.$
$\mathbf{D^2_7 B_2} : y^4 + zx + \eta x^2, yz + \xi x^2, yx + \theta x^2, z^2 + \iota x^2, zx^2, x^3.$
$\mathbf{D^3_8 B_3} : y^4 + yz + \xi x^2, y^3 z + zx + \eta x^2, yx + \theta x^2, z^2 + \iota x^2, zx^2, x^3.$
$\mathbf{D^3_8 B_4} : y^4 + yz + \xi x^2, yx + \theta x^2, z^2 + \iota x^2, zx + \eta x^2, x^3.$
$\mathbf{D^2_8 B_2} : y^4 + \eta x^2, y^2 z + (\xi + \eta) x^2, y^2 x + \theta x^2, z^2 + \iota x^2, zx^2, x^3.$
$\mathbf{D^2_8 B_3} : y^4 + y^2 z, y^2 x, z^2, x^2.$
$\mathbf{D^2_8 B_4} : y^4 + y^2 z, y^2 x, z^2, x^2.$
$\mathbf{D^1_8 B_2} : y^4 + y^2 zx + \eta x^2, y^3 z + y^2 zx + (\eta + \xi) x^2, y^3 x + \theta x^2, yx^2, z^2 + \iota x^2, zx^2, x^3.$
$\mathbf{D^1_8 B_3} : y^4 + y^3 z + \xi x^2, y^3 x + \theta x^2, y^2 zx + \eta x^2, yx^2, z^2 + \iota x^2, zx^2, x^3.$
$\mathbf{D^1_8 B_4} : y^4 + y^3 z + \xi x^2, y^3 x + \theta x^2, y^2 zx + \eta x^2, yx^2, z^2 + \iota x^2, zx^2, x^3.$
$\mathbf{D^0_8 B_2} : y^4, z^2, x^2.$
$\mathbf{D^0_8 B_3} : y^4, z^2, x^2.$
$\mathbf{D^0_8 B_4} : y^4, z^2, x^2.$
$\mathbf{D^3_9 B_3} : y^6 + zx + \eta x^2, yz + \xi x^2, yx + \theta x^2, z^2 + \iota x^2, zx^2, x^3.$
$\mathbf{D^4_{10} B_4} : y^5 + yz + \xi x^2, y^4 z + zx + \eta x^2, yx + \theta x^2, z^2 + \iota x^2, zx^2, x^3.$
$\mathbf{D^4_{10} B_5} : y^5 + yz + \xi x^2, yx + \theta x^2, z^2 + \iota x^2, zx + \eta x^2, x^3.$
$\mathbf{D^3_{10} B_3} : y^5 + y^2 z + \xi x^2, y^3 z + \xi y x^2 + \eta x^2, y^2 x + \theta x^2, z^2 + \iota x^2, zx^2, x^3.$
$\mathbf{D^3_{10} B_4} : y^5 + y^2 z, y^2 x, z^2, x^2.$

$\mathbf{D^3_{10}B_5}$: y^5+y^2z, y^2x, z^2, x^2.
$\mathbf{D^2_{10}B_3}$: $y^5+y^3z+\xi x^2, y^4z+y^2zx+\eta x^2, y^3x+\theta x^2, yx^2, z^2+\iota x^2, zx^2, x^3$.
$\mathbf{D^2_{10}B_4}$: $y^5+y^3z+\xi x^2, y^3x+\theta x^2, y^2zx+\eta x^2, yx^2, z^2+\iota x^2, zx^2, x^3$.
$\mathbf{D^2_{10}B_5}$: $y^5+y^3z+\xi x^2, y^3x+\theta x^2, y^2zx+\eta x^2, yx^2, z^2+\iota x^2, zx^2, x^3$.
$\mathbf{D^1_{10}B_3}$: y^5+y^4z, y^4x, z^2, x^2.
$\mathbf{D^1_{10}B_4}$: y^5+y^4z, y^4x, z^2, x^2.
$\mathbf{D^1_{10}B_5}$: y^5+y^4z, y^4x, z^2, x^2.
$\mathbf{D^4_{11}B_4}$: $y^8+zx+\eta x^2, yz+\xi x^2, yx+\theta x^2, z^2+\iota x^2, zx^2, x^3$.

$\vdots$

$\mathbf{D^1_5C_3}$: $x^2+xz+\iota y^2, xy+y^2, z^2, zy, y^3$.
$\mathbf{D^0_5C_3}$: x^2, z^2, y^2.
$\mathbf{D^2_6C_4}$: $x^2+xz+\eta y^4, xy+\theta y^4, z^2+\iota y^4, zy+\xi y^4+y^3, y^5$.
$\mathbf{D^1_6C_3}$: $x^2+\iota y^2, xy^2+\theta y^3, z^2, zy^2+(1+\xi)y^3, y^4$.
$\mathbf{D^1_6C_4}$: $x^2, xy^2, z^2, zy^2+y^3, y^4$.
$\mathbf{D^0_6C_3}$: $x^2+(\xi+1)xy^2+\theta zy^2+\iota y^2, z^2, y^3$.
$\mathbf{D^0_6C_4}$: x^2+xy^2, z^2, y^3.
$\mathbf{D^2_7C_5}$: $x^2+xz+\iota y^4, xy+y^3, z^2, zy, y^5$.
$\mathbf{D^1_7C_4}$: $x^2, xy^2+y^3, z^2, zy^2, y^4$.
$\mathbf{D^1_7C_5}$: $x^2, xy^2+y^3, z^2, zy^2, y^4$.
$\mathbf{D^0_7C_4}$: x^2+zy^2, z^2, y^3.
$\mathbf{D^0_7C_5}$: x^2+zy^2, z^2, y^3.
$\mathbf{D^3_8C_6}$: $x^2+xz+\eta y^6, xy+\theta y^6, z^2+\iota y^6, zy+\xi y^6+y^4, y^7$.
$\mathbf{D^2_8C_4}$: $x^2+\eta y^4, xy^2+\theta y^4, z^2+\iota y^4, zy^2+(1+\xi)y^4, y^6$.
$\mathbf{D^2_8C_5}$: $x^2+\iota y^4, xy^2+\theta y^5, z^2, zy^2+\xi y^5+y^4, y^6$.
$\mathbf{D^2_8C_6}$: $x^2, xy^2, z^2, zy^2+y^4, y^6$.
$\mathbf{D^1_8C_4}$: $x^2+xzy^2+\eta y^4, xy^3+\theta y^4, z^2+\iota y^4, zy^3+(1+\xi)y^4, y^5$.
$\mathbf{D^1_8C_5}$: $x^2+xzy^2+\iota y^4, xy^3, z^2, zy^3+y^4, y^5$.
$\mathbf{D^1_8C_6}$: $x^2+xzy^2, xy^3, z^2, zy^3+y^4, y^5$.
$\mathbf{D^0_8C_4}$: x^2, z^2, y^4.
$\mathbf{D^0_8C_5}$: x^2, z^2, y^4.
$\mathbf{D^0_8C_6}$: x^2, z^2, y^4.
$\mathbf{D^3_9C_7}$: $x^2+xz+\iota y^6, xy+y^4, z^2, zy, y^7$.
$\mathbf{D^2_9C_5}$: $x^2+\iota y^4, xy^2+\theta y^5+y^4, z^2, zy^2+\xi y^5, y^6$.
$\mathbf{D^2_9C_6}$: $x^2, xy^2+y^4, z^2, zy^2, y^6$.
$\mathbf{D^2_9C_7}$: $x^2, xy^2+y^4, z^2, zy^2, y^6$.
$\mathbf{D^1_9C_5}$: $x^2+xzy^2+\iota y^4, xy^3+y^4, z^2, zy^3, y^5$.
$\mathbf{D^1_9C_6}$: $x^2+xzy^2, xy^3+y^4, z^2, zy^3, y^5$.
$\mathbf{D^1_9C_7}$: $x^2+xzy^2, xy^3+y^4, z^2, zy^3, y^5$.
$\mathbf{D^0_9C_5}$: x^2, z^2, y^4.
$\mathbf{D^0_9C_6}$: x^2, z^2, y^4.
$\mathbf{D^0_9C_7}$: x^2, z^2, y^4.
$\mathbf{D^4_{10}C_8}$: $x^2+xz+\eta y^8, xy+\theta y^8, z^2+\iota y^8, zy+\xi y^8+y^5, y^9$.
$\mathbf{D^3_{10}C_6}$: $x^2+\eta y^6, xy^2+\theta y^6, z^2+\iota y^6, zy^2+\xi y^6+y^5, y^8$.
$\mathbf{D^3_{10}C_7}$: $x^2+\iota y^6, xy^2+\theta y^7, z^2, zy^2+\xi y^7+y^5, y^8$.
$\mathbf{D^3_{10}C_8}$: $x^2, xy^2, z^2, zy^2+y^5, y^8$.
$\mathbf{D^2_{10}C_5}$: $x^2+xzy^2+\iota y^4, xy^3+\theta y^5, z^2+\eta y^6, zy^3+(1+\xi)y^5, y^7$.
$\mathbf{D^2_{10}C_6}$: $x^2+xzy^2+\eta y^6, xy^3+\theta y^6, z^2+\iota y^6, zy^3+\xi y^6+y^5, y^7$.
$\mathbf{D^2_{10}C_7}$: $x^2+xzy^2+\iota y^6, xy^3, z^2, zy^3+y^5, y^7$.
$\mathbf{D^2_{10}C_8}$: $x^2+xzy^2, xy^3, z^2, zy^3+y^5, y^7$.
$\mathbf{D^1_{10}C_5}$: $x^2+\iota y^4, xy^4+\theta y^5, z^2, zy^4+(1+\xi)y^5, y^6$.
$\mathbf{D^1_{10}C_6}$: $x^2, xy^4, z^2, zy^4+y^5, y^6$.
$\mathbf{D^1_{10}C_7}$: $x^2, xy^4, z^2, zy^4+y^5, y^6$.
$\mathbf{D^1_{10}C_8}$: $x^2, xy^4, z^2, zy^4+y^5, y^6$.
$\mathbf{D^0_{10}C_5}$: $x^2+(\xi+1)xy^4+\theta zy^4+\iota y^4, z^2, y^5$.
$\mathbf{D^0_{10}C_6}$: x^2+xy^4, z^2, y^5.
$\mathbf{D^0_{10}C_7}$: x^2+xy^4, z^2, y^5.
$\mathbf{D^0_{10}C_8}$: x^2+xy^4, z^2, y^5.
$\mathbf{D^4_{11}C_9}$: $x^2+xz+\iota y^8, xy+y^5, z^2, zy, y^9$.
$\mathbf{D^3_{11}C_7}$: $x^2+\iota y^6, xy^2+\theta y^7+y^5, z^2, zy^2+\xi y^7, y^8$.
$\mathbf{D^3_{11}C_8}$: $x^2, xy^2+y^5, z^2, zy^2, y^8$.
$\mathbf{D^3_{11}C_9}$: $x^2, xy^2+y^5, z^2, zy^2, y^8$.
$\mathbf{D^2_{11}C_6}$: $x^2+xzy^2+\eta y^6, xy^3+\theta y^6+y^5, z^2+\iota y^6, zy^3+\xi y^6, y^7$.
$\mathbf{D^2_{11}C_7}$: $x^2+xzy^2+\iota y^6, xy^3+y^5, z^2, zy^3, y^7$.
$\mathbf{D^2_{11}C_8}$: $x^2+xzy^2, xy^3+y^5, z^2, zy^3, y^7$.
$\mathbf{D^2_{11}C_9}$: $x^2+xzy^2, xy^3+y^5, z^2, zy^3, y^7$.

$\mathbf{D}^{\mathbf{1}}_{\mathbf{11}}\mathbf{C_6} : x^2, xy^4+y^5, z^2, zy^4, y^6.$
$\mathbf{D}^{\mathbf{1}}_{\mathbf{11}}\mathbf{C_7} : x^2, xy^4+y^5, z^2, zy^4, y^6.$
$\mathbf{D}^{\mathbf{1}}_{\mathbf{11}}\mathbf{C_8} : x^2, xy^4+y^5, z^2, zy^4, y^6.$
$\mathbf{D}^{\mathbf{1}}_{\mathbf{11}}\mathbf{C_9} : x^2, xy^4+y^5, z^2, zy^4, y^6.$
$\mathbf{D}^{\mathbf{0}}_{\mathbf{11}}\mathbf{C_6} : x^2+zy^4, z^2, y^5.$
$\mathbf{D}^{\mathbf{0}}_{\mathbf{11}}\mathbf{C_7} : x^2+zy^4, z^2, y^5.$
$\mathbf{D}^{\mathbf{0}}_{\mathbf{11}}\mathbf{C_8} : x^2+zy^4, z^2, y^5.$
$\mathbf{D}^{\mathbf{0}}_{\mathbf{11}}\mathbf{C_9} : x^2+zy^4, z^2, y^5.$
$\vdots$
$\mathbf{E}^{\mathbf{3}}_{\mathbf{7}}\mathbf{F_4} : x^2+y^3+yz+\xi z^2/\iota, xy+\theta z^2/\iota, xz+\eta z^2/\iota, yz+z^2/\iota, y^4+z^2/\iota, yz^2, z^3.$
$\mathbf{E}^{\mathbf{2}}_{\mathbf{7}}\mathbf{F_4} : x^2, xy^2+y^2z, y^3, z^2.$
$\mathbf{E}^{\mathbf{1}}_{\mathbf{7}}\mathbf{F_4} : x^2+y^3, xy^2, y^4, z^2.$
$\mathbf{E}^{\mathbf{3}}_{\mathbf{8}}\mathbf{F_4} : x^2, xy^2, y^4, z^2.$
$\mathbf{E}^{\mathbf{2}}_{\mathbf{8}}\mathbf{F_4} : x^2+y^2z, xy^2, y^4, z^2.$
$\mathbf{E}^{\mathbf{1}}_{\mathbf{8}}\mathbf{F_4} : x^2+y^3z+\xi z^2/\iota, xy^3+\theta z^2/\iota, xy^2z+\eta z^2/\iota, xz^2, y^4+z^2/\iota, yz^2, z^3.$
$\mathbf{E}^{\mathbf{0}}_{\mathbf{8}}\mathbf{F_4} : x^2, y^4, z^2.$

The Tjurina numbers can be read off from these Gröbner bases. It is also checked that the associated graded ring of the Tjurina algebra over $k(\xi,\eta,\theta,\iota)$ with respect to the nilradical (x,y,z) is essentially obtained by coefficient extension of that of the closed fiber, i.e. one has the isomorphism

$$\begin{aligned}&\mathrm{gr}_{(x,y,z)}k(\xi,\eta,\theta,\iota)[x,y,z]/(\partial F/\partial x,\partial F/\partial y,\partial F/\partial z,F)\\ &\quad\cong k(\xi,\eta,\theta,\iota)\otimes_k \mathrm{gr}_{(x,y,z)}k[x,y,z]/(\partial g/\partial x,\partial g/\partial y,\partial g/\partial z,g),\end{aligned}$$

where $g(x,y,z)=0$ is the defining equation of the rational double point on the closed fiber.

4. On rationality

Whether Grauert-Riemenschneider vanishing theorem holds for our examples needs to be clarified.

Corollary 4.1. *Let X be a hypersurface singularity given by one of the equations in Theorem 3.1. Then the following assertions hold.*

i) $R^i\pi_*\mathcal{O}_{\tilde{X}}=0$ $(i>0)$ *holds for any resolution of singularities* $\pi:\tilde{X}\to X$.
ii) $R^i\pi_*K_{\tilde{X}}=0$ $(i>0)$ *holds for any resolution of singularities* $\pi:\tilde{X}\to X$.

Proof. First note that assertion i) and ii) are equivalent to each other because we have a crepant resolution $\pi:\tilde{X}\to X$, from which follows the equality $K_{\tilde{X}}\cong\mathcal{O}_{\tilde{X}}$.

From the previous theorem we know that a general hyperplane section $H\subset X$ has a rational double point. Let $\pi_1:X_1\to X$ be the blow-up along the singular locus of X. Bertini's theorem tells us that this π_1 restricted to H is also a point blow-up. The locus of a rational double point of π_1^*H corresponds to the singular locus of X_1. Then a general hyperplane section $H_1\subset X_1$ also has a rational double point, and direct calculation of cohomologies gives the vanishing. □

It might be worth mentioning that our examples are irrational from the viewpoint of the theory of tight closures.

Proposition 4.2. *Let X be a hypersurface singularity given by one of the equations in Theorem 3.1. Then the following assertions hold.*

i) *X is F-pure if and only if the type is one of* $\mathbf{E}^{\mathbf{3}}_{\mathbf{7}}\mathbf{F_4}$, $\mathbf{D}^{\mathbf{1}}_{\mathbf{4}}\mathbf{B_1}$, $\mathbf{D}^{\mathbf{1}}_{\mathbf{4}}\mathbf{B_2}$, $\mathbf{D}^{\mathbf{1}}_{\mathbf{5}}\mathbf{C_3}$, $\mathbf{D}^{\mathbf{n-1}}_{\mathbf{2n}}\mathbf{B_{n-1}}$, $\mathbf{D}^{\mathbf{n-1}}_{\mathbf{2n}}\mathbf{B_n}$, $\mathbf{D}^{\mathbf{n-1}}_{\mathbf{2n}}\mathbf{C_{2n-2}}$, $\mathbf{D}^{\mathbf{n-1}}_{\mathbf{2n+1}}\mathbf{B_{n-1}}$, $\mathbf{D}^{\mathbf{n-1}}_{\mathbf{2n+1}}\mathbf{C_{2n-1}}$ *with* $n\geq 3$.
ii) *X is not F-rational.*

Proof. i) One uses Fedder's criterion for F-purity [7, Proposition 2.1].
ii) One needs to check that the singularities which turn out to be F-pure in i) do not satisfy the criterion for F-regularity [8, Theorem 4.1.1], [9, Proposition 3.1]. Then use the equivalence of F-regularity and F-rationality for Gorenstein rings [3, §10, Proposition 10.3.7]. □

Remark 4.3. We presented here a direct proof on F-purity and F-rationality. But this result can be derived from the knowledge of types of rational double points on general hyperplane sections only. For example, combine [7, Theorem 3.4], [3, Proposition 10.3.11] and [16, Theorem 6.1]. Then as Hara points out [10, Remark 1.3], one can determine the F-purity and F-regularity of rational double points using the criteria for the defining equations given by Artin [2].

Acknowledgement

The author would like to express his sincere gratitude to Professors Kei-ichi Watanabe, Shihoko Ishii, Tadashi Tomaru, Masataka Tomari for valuable suggestions and comments. I also thank Professors Kenji Matsuki, Noboru Nakayama, Takehiko Yasuda, Shunsuke Takagi, Nobuo Hara for discussion and comments and Professors Natsuo Saito, Hiroyuki Ito and Toshiyuki Katsura for their genuine support.

References

1. M. Artin, *On isolated rational singularities of surfaces*, Amer. J. Math. **88** (1966), 129–136.
2. M. Artin, *Coverings of the rational double points in characteristic p*, in: W. L. Baily Jr. and T. Shioda (eds.), Complex Analysis and Algebraic Geometry, Cambridge Univ. Press, Cambridge, 1977, pp. 11–22.
3. W. Bruns and H. Herzog, Cohen-Macaulay Rings, Cambridge Univ. Press, Cambridge, 1998.
4. V. Cossart and O. Piltant, *Resolution of singularities of threefolds in positive characteristic.* I, J. Algebra, **320** (2008), 1051–1082.
5. V. Cossart and O. Piltant, *Resolution of singularities of threefolds in positive characteristic.* II, J. Algebra, **321** (2009), 1836–1976.
6. S. D. Cutkosky, *Resolution of singularities for* 3*-folds in positive characteristic*, Amer. J. Math. **131** (2009), 59–127.
7. R. Fedder, *F-purity and rational singularity*, Trans. Amer. Math. Soc. **278** (1983), 461–480.
8. D. Glassbrenner, *Invariant rings of group actions, determinant rings, and tight closure*, Ph. D. thesis, University of Michigan, 1992.
9. N. Hara, *F-regularity and F-purity of graded rings*, J. Algebra **172** (1995), 804–818.
10. N. Hara, *Classification of two-dimensional F-regular and F-pure singularities*, Adv. Math. **133** (1998), 33–53.
11. R. Hartshorne, Algebraic Geometry, Springer-Verlag, New York-Heidelberg, 1977.
12. M. Hirokado, H. Ito, N. Saito, *Three dimensional canonical singularities in codimension two in positive characteristic*, J. Algebra **373** (2013), 207–222.
13. J. Lipman, *Rational singularities with applications to algebraic surfaces and unique factorization*, Publ. Math. Inst. Hautes Études Sci. **9** (1969), 195–270.
14. H. Matsumura, Commutative Ring Theory, Cambridge Univ. Press, Cambridge, 1986.
15. M. Reid, *Canonical* 3*-folds*, in: A. Beauville (ed.), Journées de géométrie algébrique d'Angers 1979, Sijthoff & Noordhoff, Alphen, 1980, pp. 273–310.
16. K. Schwede and W. Zhang, *Bertini theorems for F-singularities*, Proc. London Math. Soc. (3) **107** (2013), 851–874.

Graduate school of Information Sciences, Hiroshima City University, Ozuka-higashi, Asaminami-ku, Hiroshima 731-3194, Japan

E-mail address: hirokado@math.info.hiroshima-cu.ac.jp

TABLE 4. Non-classical compound Du Val singularities in $p = 2$, I. (conti. $n \geq 4$)

	Type	Defining equation	Condition
D^r_{2n}	$\mathbf{D^r_{2n}B_r}$	$z^2 + xy^n + xy^{n-r}z + wx^2 + y^{2r+1} = 0$	$n/2<r\leq n-1$
	$\mathbf{D^r_{2n}B_{r+1}}$	$z^2 + xy^n + xy^{n-r}z + wx^2 + y^{2r+3} = 0$	
	⋮	⋮	
	$\mathbf{D^r_{2n}B_{n-1}}$	$z^2 + xy^n + xy^{n-r}z + wx^2 + y^{2n-1} = 0$	
	$\mathbf{D^r_{2n}B_n}$	$z^2 + xy^n + xy^{n-r}z + wx^2 = 0$	
	$\mathbf{D^r_{2n}C_{2r}}$	$z^2 + x^2y + xy^n + xy^{n-r}z + wy^{2r} = 0$	
	$\mathbf{D^r_{2n}C_{2r+1}}$	$z^2 + x^2y + xy^n + xy^{n-r}z + wy^{2r+1} = 0$	
	⋮	⋮	
	$\mathbf{D^r_{2n}C_{2n-2}}$	$z^2 + x^2y + xy^n + xy^{n-r}z + wy^{2n-2} = 0$	
⋮	⋮	⋮	
D^r_{2n}	$\mathbf{D^r_{2n}B_{\lfloor (n+1)/2\rfloor}}$	$z^2 + xy^n + xy^{n-r}z + wx^2 + y^{2\lfloor (n+1)/2\rfloor+1} = 0$	$1\leq r\leq n/2$
	$\mathbf{D^r_{2n}B_{\lfloor (n+1)/2\rfloor+1}}$	$z^2 + xy^n + xy^{n-r}z + wx^2 + y^{2\lfloor (n+1)/2\rfloor+3} = 0$	
	⋮	⋮	$\lfloor\ \rfloor$ stands for rounding down.
	$\mathbf{D^r_{2n}B_{n-1}}$	$z^2 + xy^n + xy^{n-r}z + wx^2 + y^{2n-1} = 0$	
	$\mathbf{D^r_{2n}B_n}$	$z^2 + xy^n + xy^{n-r}z + wx^2 = 0$	
	$\mathbf{D^r_{2n}C_n}$	$z^2 + x^2y + xy^n + xy^{n-r}z + wy^n = 0$	
	$\mathbf{D^r_{2n}C_{n+1}}$	$z^2 + x^2y + xy^n + xy^{n-r}z + wy^{n+1} = 0$	
	⋮	⋮	
	$\mathbf{D^r_{2n}C_{2n-3}}$	$z^2 + x^2y + xy^n + xy^{n-r}z + wy^{2n-3} = 0$	
	$\mathbf{D^r_{2n}C_{2n-2}}$	$z^2 + x^2y + xy^n + xy^{n-r}z + wy^{2n-2} = 0$	
⋮	⋮	⋮	
D^0_{2n}	$\mathbf{D^0_{2n}B_{n/2}}$	$z^2 + xy^n + wx^2 + y^{n+1} = 0$	n is even.
	$\mathbf{D^0_{2n}B_{(n+2)/2}}$	$z^2 + xy^n + wx^2 + y^{n+3} = 0$	n is even.
	⋮	⋮	⋮
	$\mathbf{D^0_{2n}B_{n-1}}$	$z^2 + xy^n + wx^2 + y^{2n-1} = 0$	n is even.
	$\mathbf{D^0_{2n}B_n}$	$z^2 + xy^n + wx^2 = 0$	n is even.
	$\mathbf{D^0_{2n}C_n}$	$z^2 + x^2y + xy^n + wy^n = 0$	
	$\mathbf{D^0_{2n}C_{n+1}}$	$z^2 + x^2y + xy^n + wy^{n+1} = 0$	
	⋮	⋮	
	$\mathbf{D^0_{2n}C_{2n-3}}$	$z^2 + x^2y + xy^n + wy^{2n-3} = 0$	
	$\mathbf{D^0_{2n}C_{2n-2}}$	$z^2 + x^2y + xy^n + wy^{2n-2} = 0$	
D^{n-1}_{2n+1}	$\mathbf{D^{n-1}_{2n+1}B_{n-1}}$	$z^2 + xyz + wx^2 + y^{2n-1} = 0$	
	$\mathbf{D^{n-1}_{2n+1}C_{2n-1}}$	$z^2 + x^2y + y^nz + xyz + wy^{2n-1} = 0$	
⋮	⋮	⋮	
D^r_{2n+1}	$\mathbf{D^r_{2n+1}C_{2r+1}}$	$z^2 + x^2y + xy^{n-r}z + y^nz + wy^{2r+1} = 0$	$n/2<r<n-1$
	$\mathbf{D^r_{2n+1}C_{2r+2}}$	$z^2 + x^2y + xy^{n-r}z + y^nz + wy^{2r+2} = 0$	
	⋮	⋮	
	$\mathbf{D^r_{2n+1}C_{2n-2}}$	$z^2 + x^2y + xy^{n-r}z + y^nz + wy^{2n-2} = 0$	
	$\mathbf{D^r_{2n+1}C_{2n-1}}$	$z^2 + x^2y + xy^{n-r}z + y^nz + wy^{2n-1} = 0$	
⋮	⋮	⋮	
D^r_{2n+1}	$\mathbf{D^r_{2n+1}C_{n+1}}$	$z^2 + x^2y + xy^{n-r}z + y^nz + wy^{n+1} = 0$	$1\leq r\leq n/2$
	$\mathbf{D^r_{2n+1}C_{n+2}}$	$z^2 + x^2y + xy^{n-r}z + y^nz + wy^{n+2} = 0$	
	⋮	⋮	
	$\mathbf{D^r_{2n+1}C_{2n-2}}$	$z^2 + x^2y + xy^{n-r}z + y^nz + wy^{2n-2} = 0$	
	$\mathbf{D^r_{2n+1}C_{2n-1}}$	$z^2 + x^2y + xy^{n-r}z + y^nz + wy^{2n-1} = 0$	
⋮	⋮	⋮	
D^0_{2n+1}	$\mathbf{D^0_{2n+1}C_{n+1}}$	$z^2 + x^2y + y^nz + wy^{n+1} = 0$	
	$\mathbf{D^0_{2n+1}C_{n+2}}$	$z^2 + x^2y + y^nz + wy^{n+2} = 0$	
	⋮	⋮	
	$\mathbf{D^0_{2n+1}C_{2n-2}}$	$z^2 + x^2y + y^nz + wy^{2n-2} = 0$	
	$\mathbf{D^0_{2n+1}C_{2n-1}}$	$z^2 + x^2y + y^nz + wy^{2n-1} = 0$	